

Cineraria (Dusty Miller) *maritima* Quicksilver



Life cycle	Hardy perennial
Family/origin	Asteraceae/Mediterranean region
Popular uses	Outdoor containers, pot and pack, suitable for borders, beds and silver color accents for mixed containers
Mature plant height	20-25 cm (8-10")
Mature plant width	20-25 cm (8-10")
Pot size	Packs 10-15 cm (4-6") pots
Plants per pot	1 per 10 cm (4") 3 per 15 cm (6") or larger
Sun exposure	Sun or partial shade
Water requirements	Reduced, avoid wet conditions
Flowering time	Second year flowering
Media	A well-drained, porous media is best to prevent over watering pH 5.5-6.3 EC 0.5-1.8 (50 to 150 ppm)
Growing temperature	Temperature for active growth 15°C (58°F) at night 16-18°C (60-64°F) at day time
Light	Not necessary
Fertilizer needs	Balanced N:P:K fertiliser as needed ammonium based fertilizers are not recommended, use calcium based fertilizers EC 1.8-2.2 (50-200 ppm)
Crop time	5-7 weeks
Hardiness zone	7-10



dwarf Quicksilver

Growth regulators

Quicksilver is naturally dwarf and does not need growth regulators to provide a saleable plant. However, PGRs can be applied to provide extra compact plants Daminozide (Dazide/B-Nine) Chloromequat Cycocel is possible.

Common Diseases

Botrytis, Powdery, Downy mildew and leaf spots. Avoid cool and wet conditions

Common Pests

Spider mites, Thrips, Aphids and Whitefly

Pinching

No need



Cineraria (Dusty Miller) *maritima* Quicksilver

Plug Production



Seed forms	Profi- and pelleted seed
Seed count	2.100 sds/gr (59.500 sds/oz) <i>Varies by lot!</i>
Recommended plug size	480-264
Seeds cell	2-4
Vermiculite cover	Light
Stage 1 Germination	Germination time 2-4 days Media pH 5.5 to 6.3 EC 0.5-0.75 Moisture: keep medium saturated Temperature: 20-22°C (68-72°F) growing medium temperature Fertilization: EC 0.5-1 (75-100 ppm)
Stage 2 Stem and cotyledon	Reduce moisture levels to obtain optimum root penetration and to prevent fungal diseases Prevent wet foliage after nightfall Temperature: 18-20°C (65-68°F) Fertilization: EC 0,7-1.2 (100-175 ppm)
Stage 3 Growth and development	Moisture: reduce moisture levels and allow medium to dry between watering Do not allow plants to wilt Temperature: 18-20°C (65-68°F) Fertilization: as stage 2 To prevent stretching alternate with potassium nitrate at EC 1.2-1.5 (100-175 ppm)
Stage 4 Ready for transplanting	Moisture: reduce moisture levels and allow medium to dry between watering Do not allow plants to wilt Temperature: 13-15°C (54-59°F) Fertilization: see stage 3
Plug crop time	5-6 weeks
Approximate finish time	10-12 weeks
Special tips	Ensure best light conditions especially early season and grow dry to restrict and harden growth. Transplant on time

